

Stress response and structural transitions in sheared gyroidal and lamellar amphiphilic mesophases: lattice-Boltzmann simulations

Nélido González-Segredo*

FOM Institute for Atomic and Molecular Physics (AMOLF),
P. O. Box 41883, 1009 DB Amsterdam, The Netherlands

Jens Harting†

Institut für Computerphysik, Pfaffenwaldring 27, D-70569 Stuttgart, Germany

Giovanni Giupponi and Peter V. Coveney‡

Centre for Computational Science, Department of Chemistry,
University College London, 20 Gordon Street, London WC1H 0AJ, United Kingdom

(Dated: February 6, 2008)

We report on the stress response of gyroidal and lamellar amphiphilic mesophases to steady shear simulated using a bottom-up lattice-Boltzmann model for amphiphilic fluids and sliding periodic (Lees-Edwards) boundary conditions. We study the gyroid *per se* (above the sponge-gyroid transition, of high crystallinity) and the molten gyroid (within such a transition, of shorter-range order). We find that both mesophases exhibit shear-thinning, more pronounced and at lower strain rates for the molten gyroid. At late times after the onset of shear, the skeleton of the crystalline gyroid becomes a structure of interconnected irregular tubes and toroidal rings, mostly oriented along the velocity ramp imposed by the shear, in contradistinction with free-energy Langevin-diffusion studies which yield a much simpler structure of disentangled tubes. We also compare the shear stress and deformation of lamellar mesophases with and without amphiphile when subjected to the same shear flow applied normal to the lamellae. We find that the presence of amphiphile allows (a) the shear stress at late times to be higher than in the case without amphiphile, and (b) the formation of rich patterns on the sheared interface, characterised by alternating regions of high and low curvature.

I. INTRODUCTION

The study of the response to shear in amphiphilic mesophases has been the subject of attention for numerical modellers only in recent years. The interest in the subject is sustained not only by the wide range of applications in materials' science and chemical engineering, but also by the need to gain a fundamental understanding of the universal laws governing the self-assembly processes and competing mechanisms present.

Hitherto, studies have focused mainly on the structural changes induced by steady and oscillatory shear, near and far from critical points, in polymer systems [1–6]. The morphologies studied include cubic- and wormlike-micellar, lamellar and hexagonally-packed-tubular mesophases; more complex structures are the so-called bicontinuous mesophases, of which those liquid-crystalline of cubic symmetry have thus far been considered in far less detail.

The amphiphilic gyroid [14, 15] is a bicontinuous cubic liquid crystal consisting of multi- or mono-layer sheets of self-assembled amphiphile dividing two regions, each containing phases which are mutually immiscible, e.g. aque-

ous and hydrocarbon species. These sheets or *labyrinths* form a triply periodic minimal surface (TPMS) whose unit cell is of cubic symmetry, has zero mean curvature, no two points on it are connected by a straight segment, and has no reflexion symmetries. Their skeletons, i.e. the locus bounded by the TPMS, for each immiscible phase, form double (inter-weaving), chirally symmetric 3-fold coordinated lattices. There are lyotropic [14, 15] and thermotropic transitions between the gyroid and the microemulsion mesophase, the latter being a bicontinuous mesophase of short-range order. The morphologies in the crossover regions of these transitions show shorter-range order than the gyroid's and longer-range order than the microemulsion's, reasons for which they are termed 'molten gyroids.'

Bicontinuous cubic mesophases of monoglycerides and the lipid extract from archaebacterium *Sulfolobus solfataricus* have been found at physiological conditions in cell organelles and physiological transient processes such as membrane budding, cell permeation and the digestion of fats [7]. They can also be synthesised for important applications in membrane protein crystallisation, controlled drug release and biosensors [8, 9].

The purpose of this paper is to report on the response to shear of gyroid (G), molten-gyroid (MG) and lamellar (L_α) amphiphilic mesophases simulated using a bottom-up kinetic-theoretic model for fluid flow. The model is based on a lattice-Boltzmann (LB) method, which has proved to be a modelling tool alternative to

*Email: nelido@amolf.nl

†Email: j.harting@icml.uni-stuttgart.de

‡Email: p.v.coveney@ucl.ac.uk

and more efficient and robust than sophisticated methods based on continuum equations. This LB method adheres to a *bottom-up* complexity paradigm [13] in the sense that it is simple, fully particulate and no hypotheses of desirable macroscopic behaviour are imposed on the microdynamics—yet, we have shown in the past its ability to simulate correct segregation kinetics for immiscible fluids [20] and non-equilibrium self-assembly into amphiphilic mesophases [14, 15]. Knowing that such a simple model is capable of simulating these kinetic processes from a purely bottom-up dynamics, in this paper we investigate how hydrodynamic interactions couple with self-assembly and modify the stability and morphology of the mesophases. The novelty of our work also rests in the model’s capability to reproduce morphological transitions without having to assume a macroscopic, free-energy model, used in other LB methods [3, 10] to compute the diffusive currents substantiating self-assembly.

In addition, since our method models amphiphilic molecules as point dipoles—the simplest possible particulate model for an amphiphile—the rheological features emergent from it are expected to be universal for a broad range of amphiphilic systems. Finally, most of the numerical studies measuring the stress response of complex fluids to shear reported in the literature deal with phase-segregating fluids on one side [10], and the more complicated polymeric [11] and glassy systems on the other [12]. In this respect, the present article stands somewhere in between these two.

Our paper is structured as follows. In the next section we briefly introduce the model and describe the boundary conditions for the imposition of shear. In Section III we report on simulation data and conclude that shear-thinning occurs for both G and MG mesophases, leading to a transition to a mesophase consisting of tubular and ring-like structures as the strain increases. In section IV we reveal how the presence of amphiphile in lamellar mesophases induces the formation of rich interfacial patterns surviving shear and allows higher values of stress than in lamellar mesophases without amphiphile. Finally, we provide our conclusions in Section V.

II. THE MODEL AND THE LEES-EDWARDS BOUNDARY CONDITIONS

We utilised an existing bottom-up lattice-Boltzmann (LB) model for amphiphilic fluids [14, 15], extended to simulate shear flow by means of Lees-Edwards boundary conditions [16]. The model is in turn based on an extension made to the Shan-Chen bottom-up LB model for immiscible fluids to model amphiphilic-fluid flow, and employs 25 microscopic velocities, of speeds 0, 1 and $\sqrt{2}$, in three dimensions (D3Q25 lattice) [17, 18]. The model uses a BGK (relaxation-time) approximation to the collision term of the Boltzmann equation for fluid transport,

which allows us to simulate, for large enough lattices [19], the Navier-Stokes (NS) momentum-balance equation in the bulk of each immiscible fluid species, namely “oil” (or “red”, ‘r’) and “water” (or “blue”, ‘b’). The model allows the simulation of correct phase-segregation kinetics in the absence [20] and presence [15] of a third, amphiphilic (surfactant-like, “s”) dipolar species. The model controls the inter-particle forces between r, b, and s species via coupling parameters (g_{br} , g_{bs} , g_{ss}), and transients are controlled via relaxation times for densities (τ^b , τ^r , τ^s) with an additional relaxation time for the orientation of the amphiphile dipoles (τ^d). In addition, the model simulates the nonequilibrium self-assembly and relaxation dynamics of sponge (L_3) and gyroid mesophases [14, 15]. The gyroids that it simulates show rigidity, arising from their crystalline ordering, which decreases as the concentration of amphiphile is reduced; indeed, a lyotropic transition causes the correlation length to decrease towards that of a sponge mesophase through a molten-gyroid state. This idea is central to the work we present here: we shall see that the mesophase’s crystalline ordering enhances its stress response; indeed, we find shear-thinning to occur at higher strain rates for gyroids than for sponges.

The Lees-Edwards boundary conditions (LEBC) were originally proposed by Lees and Edwards in the context of molecular dynamics simulations [16]. They showed that these boundary conditions would give rise to a desired linear, wedged velocity profile whilst avoiding the troublesome spatial inhomogeneities appearing when solid walls are used to induce the shear flow [21]. A particular realisation of the LEBC on a cartesian simulation box $[0, N_x] \times [0, N_y] \times [0, N_z]$ is established by letting the periodic images, at $N_x < x \leq 2N_x$ and $-N_x \leq x \leq 0$, move parallel to unit vectors $\pm \hat{z}$, respectively, both with speed U . The LEBC, in their original, molecular dynamics form, are expressed as a Galilean transformation on the position (x, y, z) and velocity (ξ_x, ξ_y, ξ_z) co-ordinates of a molecule, as follows

$$\begin{aligned} x' &\equiv x \bmod N_x \\ y' &\equiv y \bmod N_y \\ z' &\equiv \begin{cases} (z + \Delta_z) \bmod N_z & , x > N_x, \\ z \bmod N_z & , 0 \leq x \leq N_x, \\ (z - \Delta_z) \bmod N_z & , x < 0, \end{cases} \end{aligned} \quad (1)$$

$$\begin{aligned} \xi'_x &\equiv \xi_x \\ \xi'_y &\equiv \xi_y \\ \xi'_z &\equiv \begin{cases} \xi_z + U & , x > N_x, \\ \xi_z & , 0 \leq x \leq N_x, \\ \xi_z - U & , x < 0, \end{cases} \end{aligned} \quad (2)$$

where $\Delta_z \equiv U\Delta t$ is the image’s shift at time Δt after the onset of shear.

An implementation of the LEBC on our LB dynamics (LB-LEBC) differs from that used in molecular dynamics (MD-LEBC) in that the shift Δ_z is not in general a multiple of the lattice unit, as Wagner and Pagonabarraga

have pointed out [21], and hence an interpolation scheme is needed. This interpolation scheme streams the amphiphile dipoles $\mathbf{d}(\mathbf{x})$ and mass densities $n_k^\alpha(\mathbf{x})$ located at position $\mathbf{x}$ on the shearing wall, where $\mathbf{c}_k$ is the relevant discrete molecular velocity, $k = 1, \dots, 25$, for each (fluid and amphiphilic) species α .

In our LB-LEBC, while the spatial displacement follows Eqs. (1), the velocity shift cannot be enforced by replacing the continuum velocity component ξ_z in Eqs. (2) with the discrete microscopic speeds $\mathbf{c}_k \cdot \hat{\mathbf{z}}$, since the velocities $\mathbf{c}_k$ are constant vectors. Instead, this acceleration is enforced on the macroscopic fluid velocity around which the local-Maxwellian distributes molecules at equilibrium and towards which the BGK scheme relaxes, similarly to how immiscibility forces are implemented [15, 20]. This procedure guarantees that all accelerations in the fluid are ruled by the same BGK process, controllable via the shape of the distribution function and the relaxation-time parameter, including the acceleration due to the shearing walls.

The MD-LEBC give rise, at steady state (late times), to a shear state which is Galilean-invariant, i.e. no particular plane in the system is favoured over another. This is a *sine qua non* for any shearing method, and our method satisfies it too. As regards the unsteady, transient initial states, the MD-LEBC are unphysical since they cannot provide the molecular specificity (e.g. wall roughness) required in an atomistic approach to boundary effects, such as density layering and slip at wall. However, mesoscopic methods—LB is one of them—in general only describe low wavenumbers and frequencies, which means that, with respect to MD, (a) the atomistic detail of the shearing walls is largely coarse-grained and (b) the fluid structure and dynamics are much less sensitive to the atomistic detail of the walls. Since most boundary effects present in MD are absent in LB, the fact that the LE boundary conditions eliminate them does not pose a problem. This should be taken with a caveat: our gyroidal mesophases melt when placed in a solid box, which means that the approach to equilibrium is sensitive to momentum transfer with the walls, and therefore the LE boundary conditions do not mimic shearing a mesophase in confinement. (To our knowledge, no bottom-up simulations have ever reported mesophase self-assembly in confinement.) Rather, the LE boundary conditions in a LB model mimic shearing with walls which are far enough from the locations in the system where observables are probed such that microscopic boundary effects are absent.

Our LEBC implementation is embedded within an efficient parallel LB algorithm [22] which allows us to employ large lattices and hence reach the small Knudsen number limit where (a) regions away from interfaces satisfy the incompressible NS equation in the limit of low Mach numbers (Ma) [20], and (b) observables vary by less than 10% when the lateral lattice dimension is doubled. We

have previously found that the lattice size guaranteeing condition (b) is 128^3 for the parameters generating the mesophases investigated here [14, 15].

III. SHEARING GYROIDAL MESOPHASES

We sheared two gyroidal mesophases differing in the amount of amphiphile present and the value of the inter-amphiphile interaction coupling parameter. Each of these structures was allowed to self-assemble from homogeneous mixtures of oil, water and amphiphile using periodic boundary conditions. They have been appropriately characterised by probing direct and Fourier-space late-time snapshots of the density order parameter $\phi \equiv \rho^{\text{oil}} - \rho^{\text{water}}$; more precisely, they correspond to gyroid (cf. Fig. 5(a)) and molten gyroid mesophases, as previously reported by us [14, 15].

The common parameters used for both gyroids were oil and water densities flatly distributed in the range $0 < n^{(0)\text{b}} = n^{(0)\text{r}} < 0.7$, coupling strengths $g_{\text{br}} = 0.08$, $g_{\text{bs}} = -0.006$, relaxation times $\tau^{\text{b}} = \tau^{\text{r}} = \tau^{\text{s}} = \tau^{\text{d}} = 1$, and, for the amphiphile's dipoles, $\beta = 10$ and $d_0 = 1$.

Their differing parameters were surfactant densities, flatly distributed in the initial homogeneous mixture, in the ranges $0 < n^{(0)\text{s}} < 0.9$ for the gyroid and $0 < n^{(0)\text{s}} < 0.6$ for the molten gyroid, with coupling strengths $g_{\text{ss}} = -0.0045$ for the gyroid and $g_{\text{ss}} = -0.003$ for the molten gyroid. These values for the gyroid are 50% higher than those for the molten gyroid.

While the gyroid relaxes to a highly crystalline structure [23], the molten gyroid shows both shorter-range order and stronger temporal fluctuations than the former [15]. In order to obtain a sufficiently relaxed molten gyroid as an initial condition for the shear, we took the structure as evolved up to time step 32 500; regarding the gyroid, the time slice chosen was time step 15 000. For practical reasons, instead of letting the molten gyroid self-assemble starting from a homogeneous initial mixture, we upscaled a smaller molten gyroid, previously self-assembled using the same parameters on a 64^3 lattice [15], to a 128^3 lattice. Upscaling consisted in replicating identical copies of the system: the periodic boundary conditions used to generate the 64^3 system (a) guarantee that the density field is smooth across the replica boundaries, yet, for this same reason, (b) produce a molten gyroid with an additional, undesirable long-wavelength fluctuation whose periodicity is half the lattice size. The amplitude of this undesired long-wavelength fluctuation relaxes in time to a vanishingly small value, fact which provides us with the 128^3 mesophase we seek. We observed, however, that this relaxation takes place in less than 1 000 time steps [23], i.e. it is a fast transient which, therefore, does not affect the shear response at the late times that we are interested in. In other words, the late-time shear response is insensitive to a small perturbation

in the initial condition. This allowed us to take the up-scaled, unrelaxed structure as the initial condition for the molten gyroid.

It is worth noting that we did not require an elongated aspect ratio for the lattices along the direction parallel to the translation of the shearing walls since spatial density fluctuations were much smaller than the lattice size. This is not the case when shearing phase-segregating fluids without an amphiphilic, growth-arresting species, as has been previously reported using LB lattices of up to 128 : 128 : 512 sizes and aspect ratio [24].

A. Stress response and transients

Shear thinning is said to occur when the shear viscosity drops as the strain rate increases. For structured fluids such as those we study in this paper, the dynamic shear viscosity, η , is not expected to be a constant of the strain rate $\dot{\gamma} \equiv \frac{1}{2}(\partial_x u_z + \partial_z u_x)$ as is true of Newtonian fluids, for which,

$$P_{xz} = \pm 2\eta \dot{\gamma}, \quad \eta \neq \eta(\dot{\gamma}). \quad (3)$$

Here P_{xz} is one off-diagonal component of the pressure (or stress) tensor, and the sign, by convention, indicates that the pressure is exerted by the fluid element on the surroundings ('+') or from the latter on the former ('-'), respectively. We adhere in this paper to the second case. In our simulations, we apply the steady shear described in Section II, i.e., the shear is generated by the two image cells of the LB lattice located along the x -axis moving in opposite directions. As a consequence, $\partial_x u_z$ becomes the only non-vanishing component of the velocity gradient, which is also true for the P_{xz} component of the stress tensor (and P_{zx} , since the physical requirement that the vorticity, $W \equiv \frac{1}{2}(\partial_x u_z - \partial_z u_x)$, remains upper bounded requires the stress tensor to be symmetric).

As we have likewise done previously while computing diagonal components of the pressure tensor [14, 15, 20], here we measured P_{xz} from its definition as the sum of a kinetic term plus a virial mean-field term accounting for interactions and giving rise to non-ideal gas behaviour, namely,

$$\begin{aligned} P(\mathbf{x}) \equiv & \sum_{\alpha} \sum_k \rho_k^{\alpha}(\mathbf{x})(\mathbf{c}_k - \mathbf{u}(\mathbf{x}))(\mathbf{c}_k - \mathbf{u}(\mathbf{x})) \\ & + \frac{1}{4} \sum_{\alpha, \bar{\alpha}} g_{\alpha \bar{\alpha}} \sum_{\mathbf{x}'} \left[\psi^{\alpha}(\mathbf{x}) \psi^{\bar{\alpha}}(\mathbf{x}') + \psi^{\bar{\alpha}}(\mathbf{x}) \psi^{\alpha}(\mathbf{x}') \right] \times \\ & (\mathbf{x} - \mathbf{x}')(\mathbf{x} - \mathbf{x}'), \end{aligned} \quad (4)$$

where ψ has the form $\psi \equiv 1 - \exp[-n(\mathbf{x})]$, which saturates at high density values in order to avoid unbounded inter-particle forces whilst reproducing a meaningful equation of state [15]. Since the interaction matrix $\{g_{\alpha \bar{\alpha}}\}$ is symmetric with all diagonal elements identically zero, and

only nearest neighbour interactions are being considered, the virial term reduces to

$$\frac{1}{2} \sum_{\alpha \neq \bar{\alpha}} g_{\alpha \bar{\alpha}} \sum_k \psi^{\alpha}(\mathbf{x}) \psi^{\bar{\alpha}}(\mathbf{x} + \mathbf{c}_k) \mathbf{c}_k \mathbf{c}_k. \quad (5)$$

In the incompressible, low Mach number limit, our LB model reproduces the NS equation away from interfaces [17, 19], which describes a Newtonian fluid with a viscosity being a well known function of the relaxation time. The presence of an interface, characterised by an interfacial tension and a bending rigidity, however, introduces anisotropies in the fluid's stress tensor which can be accounted for by a tensorial effective viscosity. Since the interface may move, at a speed growing with the strain rate, these anisotropies can become unsteady. Our aim is then to measure how this viscosity evolves with the strain and the strain rate.

In order to probe the function $\eta = \eta(\dot{\gamma})$ for both gyroidal mesophases, we measured P_{xz} for a number of different applied shear rates. The chosen values for U were such that they remained within the incompressibility limit, i.e. small compared to the speed of sound on the D3Q25 lattice, $c_s = 3^{-1/2} \approx 0.58$. Values chosen were $U = 0.05, 0.10, 0.15, 0.20$, corresponding to Mach numbers $\text{Ma} \equiv U/c_s = 0.086, 0.17, 0.26, 0.34$, respectively. All observables we report in this paper are spatial averages, at least on $x = \text{const.}$ planes where a simple fluid under the same shear would show translational symmetry for the velocity field, i.e., perpendicular to the velocity gradient. Since, for reasons of computational cost, we do not perform averages over the seed used to generate the pseudo-random initial configuration mimicking a homogeneous ternary mixture, we do not provide error bars around averages.

Figure 1 shows the profile of the stress, for the sheared gyroid, along the applied velocity gradient direction. Several curves therein depict the transport of momentum towards the core (i.e., the plane $x = 64$) as the strain grows as a function of time. Distinctively, the profiles have spatial fluctuations, a consequence of the gyroid's convoluted structure whose interfacial tension locally modifies the viscosity expected for a simple fluid. The u_z component of the velocity field, shown in Fig. 2 and averaged in the same way as stated for $\langle -P_{xz} \rangle$ in the caption of Fig. 1, is however not inhomogeneous but follows a transient similar to that expected for a simple fluid: we observe the setting up of a steady, smooth and wedge-shaped profile, except at the borders. Figure 2 also includes the behaviour of the averaged velocity profile for the molten gyroid MG at late times, and is seen to match that of the gyroid G.

Remaining with the G mesophase, we show in Fig. 3 the temporal evolution of the stress displayed in Fig. 1; the values plotted are averages of the latter on the $8 \leq x \leq N_x - 8 = 120$ interval, which amounts to averaging over the whole lattice except thin slabs adjacent to the bound-

aries. In addition to Fig. 1, we include higher and lower shear velocities, namely $U = 0.05, 0.15, 0.20$. Note that the time evolution of the averaged stress is a succession of peaks and troughs, denoting successive intervals of yield and recoil, which is a canonical feature of viscoelastic behaviour. Were the strain rate at which the gyroid deforms coincident with the applied shear rate, these curves would imply shear thinning. In fact, while the increments in applied shear rate between these curves are kept constant, the increments in the (absolute) values of the stress at late times do not remain so but decrease. In Fig. 4 we show the stress averaged over time steps 24 000 to 28 000, plotted against the true strain rate, where the latter was measured from the linear velocity profile generated at $\Delta t \geq 9\,000$ ($t \geq 24\,000$), as displayed in Fig. 2. Figure 4 clearly shows shear thinning: the slope, i.e., the effective viscosity $\eta^{\text{eff}} \equiv \partial P_{xz} / \partial \dot{\gamma}$, decreases with the strain rate. Figure 4 also contains the analogous curve for the molten gyroid, which shows shear thinning for the latter at lower strain rates than those at which the gyroid does, and at higher intensity, i.e.

$$\left. \frac{\partial \eta^{\text{eff}}}{\partial \dot{\gamma}} \right|_{\text{molten}} < \left. \frac{\partial \eta^{\text{eff}}}{\partial \dot{\gamma}} \right|_{\text{gyroid}} < 0. \quad (6)$$

This is the first indication of shear thinning reported by means of a bottom-up kinetic-theoretic model for fluid flow.

B. Morphological transitions

Figure 5 shows the configuration of the gyroid in the $40 \leq y \leq 52$ slab of the $128 : 128 : 128$ lattice, before and at late times after applying a shear of $U = 0.20$. The volume-rendering graphical representation employed [25] makes regions where $\phi \geq 0.37$ opaque to the lighting rays, assumed to shine normal to the plane of the text and inwards; since $-0.79 \leq \phi \leq 0.79$ over the entire system, these regions are the high-density locus of one of the species (say, oil). Before shear, the structure contains highly ordered subvolumes of gyroid symmetry and diagonal length from about 32 to 64 lattice sites, cf. Fig. 5(a). This gyroid is hence a collection of subvolumes with a regular tubular structure making up two three-fold coordinated, interweaving chiral lattices of which we depict only one. Since the size of the G unit cell is approximately 5 to 6 lattice units, the depth (y -dimension) of the slabs shown in Fig. 5 is of about two gyroid unit cells. As can be seen in Fig. 5(a), the interfaces between these gyroid subvolumes are defective regions where long-range order and symmetry appears to be drastically reduced [14, 15]. Two features characterising them are the spatial variation in coordination number and chirality, seen by the presence of elongated tubules and toroidal rings, cf. figure 6.

At $\Delta t = 21\,000$, which is a late time after the onset of shear and we take as steady state, the structure has

lost any resemblance with the initial gyroid, except for the persistence of the toroidal rings, see Fig. 5(c), which are defects in G. Also, the structure at $\Delta t = 21\,000$ is essentially the same as that at time step $\Delta t = 5\,000$ —it is a nonequilibrium steady state for at least the previous 16 000 time steps, a time longer than that required for the initial configuration to self-assemble from a homogeneous mixture of oil, water and amphiphile. The structure at $\Delta t = 21\,000$ consists of an irregular network of mainly the same structural elements characterising the defective regions before the onset of shear, namely, (a) elongated tubules, with a tendency to align along a direction which is a linear combination of directions $(1, 0, 0)$ and $(0, 0, 1)$, and (b) toroidal, ring-like structures. This description is, by visual inspection, similar for every subvolume of the lattice visualised.

We also looked into the structure of the sheared molten gyroid at late times. In contradistinction to the gyroid's state at high strain, showing tubules of shape similar to that depicted in Fig. 6 and at an angle with the $x = \text{const.}$ planes, the highly strained molten gyroid displays tubes which are more stretched and aligned along the $\hat{z}$ direction. The toroidal rings, also present for the molten gyroid before shear, represent a much smaller volume fraction for the sheared molten gyroid than for the sheared gyroid.

Figure 7 shows the summed structure function $\sum_{k_y} S(\mathbf{k}, t)$, or scattering pattern, of the sheared gyroid mesophase, showing stages of its plastic deformation. Here, $S(\mathbf{k}, t)$ is the structure function, computed according to [15, 20]

$$S(\mathbf{k}, t) \equiv \frac{\varsigma}{V} \left| \phi'_{\mathbf{k}}(t) \right|^2. \quad (7)$$

Here, $\mathbf{k}$ is the discrete wavevector, V is the lattice volume, ς is the unit cell volume for the D3Q25 lattice, and $\phi'_{\mathbf{k}}(t)$ is the Fourier transform of the fluctuations of ϕ . $S(\mathbf{k}, t)$ is the Fourier transform of the autocorrelation function for the order parameter,

$$C_{\phi\phi}(\mathbf{r}, t) \equiv \langle \phi((\mathbf{x}, t)) \phi(\mathbf{x} + \mathbf{r}, t) \rangle \quad (8)$$

where $\mathbf{r}$ is a vector lag and the brackets indicate an average over the spatial coordinate $\mathbf{x}$. Figures 7(a), (b) and (d) are the xz ‘scattering patterns’ of the structures in Fig. 5, produced by summing up the structure function along the x direction. At $\Delta t = 1\,000$ (not shown), the maximum intensity is reduced to 29% of its value at $\Delta t = 0$, while there appear horizontal ‘smeared out filaments’ of very weak intensity, intrinsically related to the shearing process, as we shall conclude from Fig. 8. At $\Delta t = 5\,000$ a clear *cardioid* shape has developed; the fact that it persists for the rest of the simulation confirms our observation that the system reaches a steady state at time step $\Delta t = 5\,000$. In addition, there is no trace of gyroidal patterns along the x -direction.

In order to investigate the origin of the cardioid shape, we computed the scattering pattern for a ‘synthetic gyroid’,

$$G(\mathbf{x}) \equiv \sin qx \cos qy + \sin qy \cos(qz - \delta(\mathbf{x})) + \sin(qz - \delta(\mathbf{x})) \cos qx. \quad (9)$$

where $\delta(\mathbf{x}) = (x - N_x/2)\delta_{\max}$ is a spatially-varying dephase used to obtain a linear strain on the morphology (its maximum value, $\delta_{\max}$, is reached at the lattice boundaries), and $q = \text{const.}$ is a wavenumber controlling the size of the surface’s unit cell. It is known that $G(\mathbf{x}) = 0$ for $\delta_{\max} \equiv 0$ is a good approximation to the Schoen “G” triply periodic minimal surface of $Ia\bar{3}d$ cubic symmetry, referred to as ‘the ideal gyroid’ hereafter [26]. Figure 8 shows the scattering patterns for the unstrained morphology and for dephases $\delta_{\max} = 8, 16$.

Comparing structure function maps in Figs. 7 and 8, at the same value of the strain rate, proves useful. For the synthetic gyroid, the strain is controlled by the number of unit cells that the dephase causes the structure to shift at the lattice boundary, following a linear profile as we approach the other boundary going through zero strain at the lattice core. For our simulated amphiphilic gyroid, however, the strain does not follow a linear profile at early times; instead, the strain at time t would need to be computed from the integral $\frac{1}{N_x} \int_0^t \int_0^{N_x} dt' dx \partial_x u_z(\mathbf{x}, t')$, where t' is the time parameter. For the purposes of this paper, however, such an analysis would be superfluous; in fact, Fig. 8 already provides us with enough information to understand the origin of the cardioid shape. For all panels, (a), (b) and (c) therein, the position of the peaks at $k_z = 0$ ($k_x/(2\pi/N) \approx -14, 15$, where $N = 128$) are invariant under the strain (dephase); not so with the peaks at $k_z \neq 0$, which shift leftwards. (The shift would be rightwards were $\partial_x u_z < 0$ or $\delta_{\max}$.) The shape of the maps in Figs. 7(c) and 7(d) is that of a transformed scattering pattern shifted leftwards. This transformation occurs early, between $\Delta t = 0$ and $\Delta t = 3000$, and is characterised by two (strong, $S \geq 700$) peaks similar to those of the gyroid at $k_x = 0$, and two (weaker, $200 \leq S < 700$) peaks at $k_z = 0$.

IV. A SIMPLER CASE: SHEARING THE LAMELLAR MESOPHASE

In the last section we reported on the gyroid displaying higher shear stress than the molten gyroid. Since the structural transition between these two mesophases can be driven by both the amphiphile density and the inter-amphiphile coupling parameter, as we have reported in the past [15], our aim in this section is to elucidate the role of the amphiphile density alone on the stress response to shear; we choose the lamellar mesophase as the subject of study, since this is the mesophase with the simplest possible internal interface.

The initial configuration employed was a cubic 128^3 lattice with 16 lamellae, stacked perpendicularly to unit vector $\hat{\mathbf{z}}$. The lamellae were of alternating, oil-water compositions, separated by a thin monolayer of amphiphile; the thickness of the immiscible and amphiphilic lamellae were 7 and 1 lattice sites, respectively. We populated each lattice site with a value of density kept constant over the region corresponding to a given species; each microscopic velocity is assigned the same fraction of this value. We gave amphiphilic regions the densities $n^{(0)s} = 0, 0.80, 0.95$, and oil and water regions the densities $n^{(0)r} = n^{(0)b} = 0.7$. Shear was applied perpendicular to the lamellae with the same LBC employed in the last section, with speed $U = 0.10$.

Before the onset of shear, the case without amphiphile for the lamellar initial condition just described is, *a priori*, a metastable state in our LB model. In fact, the structure has a stationary morphology since short-range oil-water forces and the absence of fluctuations maintain immiscibility, i.e. a value for the interface steepness, $|\nabla\phi|$; however, a large enough perturbation in ϕ may allow a fluctuation in surface tension which drives the entire interface to a radically different shape. Another factor disrupting this lamellar morphology is shear, which may work against the interfacial tension by reducing $|\nabla\phi|$; this can lead to miscibility ($\phi \equiv 0$) for high enough strain rates. Despite these arguments, we observed stability for the sheared lamellar mesophase without amphiphile, as we report next.

Figure 9 shows the stress as measured in the same fashion performed on the data plotted in Fig. 3, for several amphiphile densities. The behaviour observed is diverse. For zero amphiphile concentration (solid curve), the stress reaches a peak at early times before it proceeds to a second, lower maximum at late times, going through a trough at intermediate times due to the fact that $|\nabla\phi|$ experiences a transient decrease.

The high-density regions of one of the immiscible species (say, oil) is shown in Fig. 10(a) at late times, $\Delta t = 8000$; these are representative of the shape of the oil-water interface. Away from the boundaries ($x = 0, 128$), there is a large interfacial area with zero curvature, where we define the curvature as $H \equiv \partial_{zz}^2 x_\phi(z)$, $x_\phi(z)$ being the curve resulting from projecting the $\phi = 0.18$ surface onto the xz plane. Curiously, we observe three changes of curvature as we follow the curve $x_\phi(z)$ for $y = \text{const.}$, namely, $H < 0, H > 0, H < 0, H > 0$; instead, we would have expected the steady, late-time configuration for the sheared lamellar mesophase to rather minimise the interfacial area, leaving only one inflexion point. For fluid regimes under conditions of local thermodynamic equilibrium, we can think of H^2 as an interfacial free energy density associated with the curvature [27]; in this case, we would have expected the steady, late-time configuration to also minimise the interfacial free energy.

The stress curve corresponding to $n^{(0)s} = 0.80$

(cf. Fig. 9) shows the absence of large troughs, as occurs for the $n^{(0)s} = 0$ case, despite the fact that interfacial tension is drastically reduced by the presence of the amphiphile. In addition, the stress grows at late times to higher values than those achieved by the $n^{(0)s} = 0$ case. The late-time order-parameter configuration is displayed in Fig. 10(b), showing a rich interfacial pattern. Using the same arguments as those of the last paragraph, this structure could be characterised by a higher curvature energy, $\int d^2x H^2$, where d^2x is a measure on the oil-water interface, and H is now defined as the inverse radius of curvature, parametrised on the arclength, s . Figure 10(b) shows similar regions of high curvature at an equal distance from the shearing walls, where $u_z = \text{const.}$, which we shall call *nodal* planes. Also note that the interface, as approximately depicted by the boundary of the $\phi \geq 0.22$ volume, joins the lattice boundary at an angle close to 90 degrees.

The stress curve for the $n^{(0)s} = 0.95$ case shows a dramatically different situation for the first 5 000 time steps: the presence of a trough, deeper than that present for the $n^{(0)s} = 0$ density. After that, there appears a shoot-off whereby the stress rapidly grows and equals the late time value achieved in the $n^{(0)s} = 0.80$ case, while the order-parameter displays a configuration analogous to the $n^{(0)s} = 0.80$ case, cf. Fig. 10(c). By looking at the amphiphile density field, $\rho^s(\mathbf{x})$, for the case $n^{(0)s} = 0.95$, we observed that the high curvature regions arise close to the boundaries first ($\Delta t < 1000$), and then rapidly move away from them as the strain progresses.

V. CONCLUSIONS

In this paper we have reported on the shear stress response of two gyroidal cubic amphiphilic mesophases previously self-assembled using the same bottom-up LB model we employ here, namely, the gyroid *per se*, G, which shows high crystallinity at late self-assembly times, and the molten gyroid, MG, endowed with shorter-range order and located within the sponge-gyroid lyotropic structural transition [15]. Shear was imposed via sliding periodic (Lees-Edwards) boundary conditions, and we investigated the response to several values of the strain rate. In addition, in order to investigate the dependence of the shear stress on the amphiphile density, we also sheared a lamellar mesophase, of much simpler morphology than the gyroidal mesophases.

We found that the gyroidal mesophases exhibit shear thinning, more pronounced and at lower strain rates for the MG mesophase than the G mesophase. In other words, momentum introduced into the system due to shear is transported more easily for the mesophase containing more amphiphile, with longer-range ordering, i.e. the effective viscosity is higher for the G mesophase.

We also found a shear-induced transition from an initial

gyroidal morphology (G and MG) to a mesophase characterised by coexisting elongated tubules and toroidal, ring-like structures. The features of this mesophase is in contrast to those of the mesophase reported by Zvelindovsky et al. using free-energy Langevin-diffusion methods by shearing a bicontinuous structure reminiscent of a molten gyroid [3]. The structure they found is of a shorter-range ordering than that of the MG mesophase described here, and the high-strain structure consists of coexisting lamellae and hexagonally packed tubes elongated along the direction of the imposed shear velocity. Our sheared mesophases also show elongated tubes along this direction, but the structure is far more complicated than that found by Zvelindovsky et al. in that it exhibits remnant toroidal rings and ‘hard shoulders’ reminiscent of gyroidal skeletons; hexagonal packing and coexisting lamellae are, on the other hand, absent.

The shear performs a plastic deformation which effectively breaks the links of the gyroidal skeleton; this happens as these links interpose an (oil-water) interface whose normal, $\mathbf{n}$, is parallel or anti-parallel to the flow streamlines, $\mathbf{u}$. In other words, shear effectively applies a ‘mixing’ force which is in competition with the inter-particle forces keeping the mesophase in place, namely, those controlled by coupling parameters g_{br} , g_{bs} and g_{ss} . Our hypothesis is that adsorbed dipoles sitting on interfacial regions at an angle $\theta \equiv \angle(\mathbf{u}, \mathbf{n})$ other than $\theta = 0, \pi$ require more work from the shear forces to be drawn away from the interface than those regions on which the streamline impinges normally, since the mixing force goes as $\cos \theta$. In particular, since the mixing force vanishes for $\theta = \pi/2$, considerably longer interfaces can survive the flow—shear induces a preferential direction along which the long-range order present before the onset of shear is not reduced. This explains not only the formation of the elongated tubules but also their reconnection (increase in coordination number). In fact, the toroidal, ring-like structures are not only vestigial gyroid defects which have survived the gradient $\nabla \mathbf{u}$, but are also born anew as a result from reconnections. It is relevant to point out that Padding and Boek, using a coarse-grained molecular dynamics model for wormlike micelles, reported on the formation of rings when applying steady shear to a wormlike micellar mesophase [6]—this is an ‘amphiphile-in-water’ binary mesophase, in contrast to the ‘oil-amphiphile-water’ ternary mesophases we study in this paper.

By applying shear to a lamellar mesophase we found that the presence of amphiphile on the oil-water interface of the mesophase causes the interface to fold into a wealth of structures with a (discrete) translational symmetry on planes equidistant to the shearing walls and along the direction of the shear velocity. In other words, the inter-amphiphile force couples the adsorbed amphiphilic dipoles so that the interface locally increases its curvature energy density. It is worth investigating whether this

local increase is due to the amphiphile being incapable of sustaining interfacial regions of low curvature under shear, i.e. whether the ‘breaking’ mechanism induced by shear is counteracted by regions of high curvature energy density. Regarding the shear stress, our amphiphile-containing lamellae responded with higher stress at late times than those without amphiphile. This contrasts with the results found for the gyroidal mesophases, and lets us conclude that it is the gyroid’s cubic morphology that allows this structure to be stiffer. Understanding the behaviour of the lamellar mesophase under shear requires the study of amphiphile self-assembly under shear, including in and out of plane amphiphilic and associated Marangoni currents, and their coupling to the imposed flow.

VI. ACKNOWLEDGMENTS

We thank Dr. Rafael Delgado-Buscalioni, Prof. Antonio Coniglio and Prof. Francesco Sciortino for enlight-

ening discussions. We acknowledge Iain Murray, Elena Breitmoser and Jonathan Chin for their involvement in algorithm implementation and optimisation within the RealityGrid and TeraGyroid projects. This work was supported by the UK EPSRC under grant RealityGrid GR/R67699 which also provided access to a 512-processor SGI Origin3800 platform at Computer Services for Academic Research (CSAR), Manchester Computing, UK. We also thank the Higher Education Funding Council for England (HEFCE) for our on-site 16-node SGI Onyx2 graphical supercomputer.

-
- [1] A. V. M. Zvelindovsky, B. A. C. van Vlimmeren, G. J. A. Sevink, N. M. Maurits, and J. G. E. M. Fraaije, *J. Chem. Phys.* **109**, 8751 (1998).
 - [2] A. V. M. Zvelindovsky, G. J. A. Sevink, B. A. C. van Vlimmeren, N. M. Maurits, and J. G. E. M. Fraaije, *Phys. Rev. E* **57**, R4879 (1998).
 - [3] A. V. M. Zvelindovsky, G. J. A. Sevink, and J. G. E. M. Fraaije, *Phys. Rev. E* **62**, R3063 (2000).
 - [4] A. N. Morozov, A. V. M. Zvelindovsky, and J. G. E. M. Fraaije, *Phys. Rev. E* **64**, 051 803 (2001).
 - [5] A. N. Morozov and J. G. E. M. Fraaije, *Phys. Rev. E* **65**, 031 803 (2002).
 - [6] J. T. Padding and E. S. Boek, *Phys. Rev. E* **70**, 031 502 (2004).
 - [7] P. Mariani, V. Luzzati and H. Delacroix, *J. Mol. Biol.* **204**, 165 (1988).
 - [8] S.-J. Marrink and D. P. Tieleman, *J. Am. Chem. Soc.* **123**, 12383 (2001).
 - [9] V. Luzzati, R. Vargas, P. Mariani, A. Gulik, and H. Delacroix, *J. Mol. Biol.* **229**, 540 (1993).
 - [10] A. Xu, G. Gonella, and A. Lamura, *Phys. Rev. E* **67**, 056 105 (2003).
 - [11] R. D. Groot and W. G. M. Agterof, *Macromolecules* **28**, 6284 (1995).
 - [12] P. Sollich, *Phys. Rev. E* **58**, 738 (1998).
 - [13] P. V. Coveney, *Phil. Trans. R. Soc. Lond. A* **361**, 1057 (2003).
 - [14] N. González-Segredo and P. V. Coveney, *Europhys. Lett.* **65**, 795 (2004).
 - [15] N. González-Segredo and P. V. Coveney, *Phys. Rev. E* **69**, 061501 (2004); *Virt. J. Nanoscale Sci. & Tech.* **9**, (Issue 23, June 14 2004), *Supramolecular and Biochemical Assembly*.
 - [16] A. W. Lees and S. F. Edwards, *J. Phys. C* **5**, 1921 (1972).
 - [17] H. Chen, B. M. Boghosian, P. V. Coveney, and M. Nekovee, *Proc. R. Soc. London A* **456**, 2043 (2000).
 - [18] An example of the use of the D3Q25 lattice in lattice-gas methods is: P. J. Love, P. V. Coveney, and B. M. Boghosian, *Phys. Rev. E* **64**, 021503 (2001).
 - [19] S. Succi, “The lattice-Boltzmann equation—for fluid dynamics and beyond.” (Oxford University Press, Oxford, 2001.)
 - [20] N. González-Segredo, M. Nekovee, and P. V. Coveney, *Phys. Rev. E* **67**, 046304 (2003).
 - [21] A. J. Wagner and I. Pagonabarraga, *J. Stat. Phys.* **107**, 521 (2002).
 - [22] P. J. Love, M. Nekovee, P. V. Coveney, J. Chin, N. González-Segredo, and J. M. R. Martin, *Comp. Phys. Commun.* **153**(3), 340 (2003).
 - [23] J. Harting, M. J. Harvey, J. Chin and P. V. Coveney, *Comp. Phys. Commun.* **165**, 97 (2005); J. Harting, J. Chin, M. Venturoli, P. V. Coveney, “Large-scale lattice Boltzmann simulations of complex fluids: advances through the advent of computational grids.” *Phil. Trans. R. Soc. London Series A*, in press (2005).
 - [24] J. Harting, M. Venturoli and P. V. Coveney, *Phil. Trans. R. Soc. London Series A* **362**, 1703 (2004).
 - [25] For an explanation of volume rendering, see, e.g., W. Schroeder, K. Martin, B. Lorensen, “The Visualization Toolkit: An Object Oriented Approach to 3D Graphics.” Kitware Inc., ISBN 1930934076, 3rd Edition, 2003; URL: <http://public.kitware.com/VTK/>.
 - [26] M. Wohlgenuth, N. Yufa, J. Hoffman and E. L. Thomas, *Macromolecules* **34**, 6083 (2001).
 - [27] R. Lipowsky, *Nature* **349**, 475 (1991).




FIG. 1: Shear stress response of a gyroid mesophase along the direction of the velocity gradient. As initial condition, we have taken a gyroid on a $N_x N_y N_z = 128^3$ cubic lattice at time step $t = 15\,000$ of self-assembly [14, 15]. The Lees-Edwards walls move with speed $U = 0.10$ ($\text{Ma} = 0.17$). For each x coordinate, the original field has been averaged on the plane $[1, N_y] \times [16, N_z - 16]$, where the excluded interval on the z -axis accounts for wrapped-round densities. Standard errors of the averages are about 6×10^{-8} throughout, and are not shown. Each line represents the response at Δt time steps after the start of steady shear: $\Delta t = 0$ (dotted line), $\Delta t = 100$ (dash-dotted), $\Delta t = 800$ (dashed) and $\Delta t = 9\,000$ (solid), where the last is ca. the time at which the core (i.e., the plane $x = 64$) fully responds. From the figure we can see that momentum transfer decreases as it reaches the core from the walls. Also, note that the stress inverts its sign at late times adjacent to the boundaries, $|x - x_0| \leq 2$ ($x_0 = 0, 128$). All quantities reported are in lattice units.




FIG. 2: Spatially averaged velocity component u_z for the molten gyroid and the gyroid mesophases sheared with $U = 0.10$, at late times and over the $x \geq 64$ half of the system. The dashed thin and thick curves correspond to the molten gyroid at time steps $\Delta t = 9000$ and 13000 , respectively. The solid thin and thick curves correspond to the gyroid at time steps $\Delta t = 9000$ and 13000 , respectively. The average is over the same two-dimensional domain as described in Fig. 1, for each x , and its standard error is shown as negligible error bars. Note that the velocity shows a maximum located from 2 to 4 sites away from the boundary, unlike a simple fluid which would display it exactly at the boundary. The value of this maximum coincides with the actual velocity at which the BGK relaxation process of our LB model is forcing the fluid to move, which needs not coincide with the input parameter $U = 0.10$. Note that the inversion in the sign of the stress that we reported in Fig. 1 occurs precisely for $|x - x_0| \leq 2$, $x_0 = 0, 128$ and at (late) times close to and after $\Delta t = 9000$. The behaviour at the other boundary region is similar and symmetric to that displayed here. All quantities reported are in lattice units.




FIG. 3: Temporal evolution of the average shear stress of the gyroid for different values of steady shear. The initial condition is the same as that mentioned in Fig. 1. The curves, as seen, e.g., at $\Delta t = 4000$ from bottom to top, correspond to Lees-Edwards walls moving with speeds $U = 0.05, 0.10, 0.15, 0.20$ (shear rates $S/10^{-3} = 0.39, 0.78, 1.17, 1.56$), respectively. The dotted curves are the responses after a sudden termination of shear; they are also referred to as the system's relaxation functions for the relevant shear speeds. The average here is in the three-dimensional domain $[8, N_x - 8] \times [1, N_y] \times [16, N_z - 16]$, where $N_x = N_y = N_z = 128$ and error bars are the standard error of the average. All quantities reported are in lattice units.




FIG. 4: Both the gyroid (solid line) and the molten gyroid (dashed) mesophases exhibit shear thinning. Shown is the stress averaged over the interval $24\,000 \leq t \leq 28\,000$. From the figure it is clear that the gyroid manifests greater stiffness than the molten gyroid and its (effective) viscosity drops for higher strain rates. All quantities reported are in lattice units.




FIG. 5: High-density locus of one species (say, oil) in the gyroid mesophase, before shearing, (a), and at an early, (b), and late time slice, (c), after the onset of shear. The shear speed is $U = 0.20$. The complementary immiscible fluid (water) fills the voids with a similar, inter-weaving structure. The system is on a $128 \times 128 \times 128$ lattice, and all figures show the subvolume $40 \leq y \leq 52$ and the reference system in use (the y -axis is perpendicular to the plane of the page). The initial configuration, (a), is a gyroid at 15 000 time steps of self-assembly under periodic boundary conditions. These images are volume renderings of the density order parameter, $\phi \equiv \rho^{\text{oil}} - \rho^{\text{water}}$; the regions visible to the reader are those for which $\phi \geq 0.36$ whilst over the entire fluid $-0.79 \leq \phi \leq 0.79$; the regions for which $\phi \leq -0.36$ (water, not shown) display a similar structure which is complementary (interweaving) to the one shown here. All quantities reported are in lattice units.




FIG. 6: Schematic representation of the skeleton (locus of highest density) of the gyroid mesophase we employ, and two of its structural features before and at late times after the onset of steady shear. The thickness provides a sense of perspective, and represents how close each segment is to the reader; note that the figures on the right are planar. The skeleton denoted by ‘gyroid’ depicts a portion of one of the two chiral lattices making up the long-range order regions of the gyroid before shear, cf. Fig. 5a—the coordination number is three at each node. In the regions of the gyroid containing defects, as well as in most of the sheared mesophase at late times, the coordination number can be reduced to two, describing a ‘tubule’. We also show the skeleton of the ‘ring’ structure ubiquitous in the sheared gyroid at late times, also present in smaller proportion as a defect in the mesophase before the onset of shear. At lower values of density, this ring appears to be toroidal.




FIG. 7: Projected structure function (‘scattering pattern’) as a function of the time step for the sheared gyroid, as calculated using Eq. (7). Shear velocity is $U = 0.20$. (a), (b) and (c) are scattering patterns before shear and at intermediate and late times after the onset of shear, respectively, while, for completeness, (d) details the side view of the structure function corresponding to (c). The initial condition for shearing was a gyroid on a 128^3 lattice at 15 000 time steps of self-assembling. Time steps after the start of shear for these snapshots are indicated below each. Darkness in the greyscale grows with the scattering intensity—filled isocurves correspond to values $S = 1, 80, 200, 700$. The spikes are shear-dependent features; see Fig. 8 and text for discussion. All quantities reported are in lattice units, and $N \equiv N_x = N_z$.




FIG. 8: Structure function of the ‘synthetic gyroid’, as calculated using Eq. (7) on the field $G(\mathbf{x})$, the expression of the latter being Eq. (9). Parameter $\delta_{\max}$ is the maximum value of the dephase $\delta(\mathbf{x}) = (x - N_x/2)\delta_{\max}$, which serves to mimic a uniform strain across the structure. The case $\delta_{\max} = 0$ gives an approximation to the Schoen G (or ‘ideal gyroid’) structure. Darkness in the greyscale grows with the scattering intensity, and the filled isocurves shown correspond to $S = 1, 80, 200, 700$. For $k_z \neq 0$, the strain shifts the pattern leftwards and smear the peaks, while leaving the $k_z = 0$ peaks intact. The smearing not being in direct relation to the strain—panel (b) shows more smearing than panel (c)—suggests a similar behaviour for the spikes shown in Fig. 7(c). The ‘cardioid’ shape reported in Fig. 7 originates from the facts that the structure undergoes a structural transition (weakening and/or relocation of some of its $k_z \neq 0$, $k_x \neq 0$ peaks) whilst being sheared with a velocity profile of positive slope (cf. Fig. 2), which orients the ‘atria’ leftwards. All quantities reported are in lattice units, and $N \equiv N_x = N_z$.




FIG. 9: Temporal evolution of the average shear stress response of a lamellar mesophase at a shear speed of $U = 0.10$, for different initial amphiphile densities $n^{(0)s}$. The solid, dashed, and dash-dotted curves correspond, respectively, to $n^{(0)s} = 0, 0.80, 0.95$. The average is computed over the three-dimensional domain $[8, N_x - 8] \times [1, N_y] \times [16, N_z - 16]$, where $N_x = N_y = N_z = 128$ and error bars are not included since they are negligible. All quantities reported are in lattice units.




FIG. 10: Slabs $0 \leq y \leq 8$ of the order parameter ϕ for sheared lamellar mesophases corresponding to increasing amphiphile density, $n^{(0)s}$, as indicated below each relevant panel, at time step $\Delta t = 8000$ after the onset of shear and for shear velocity $U = 0.10$. The coordinate system is the same as that in Fig. 5. In panel (a), the regions opaque to incident (volume rendering) light are those for which $\phi \geq 0.18$, where $|\phi| \leq 0.36$ across the system. In panel (b), the opaque regions are those for which $\phi \geq 0.22$, where $|\phi| \leq 0.45$ across the system. In panel (c), the opaque regions are those for which $\phi \geq 0.24$, where $|\phi| \leq 0.48$ across the system. It is worth noting that the surfactantless case, (a), exhibits a curved interface. The amphiphilic cases, (b) and (c), display the formation of irregularities in the interface and nodal planes, as a result of the inter-amphiphile interaction. All configurations have translational symmetry along the y -axis. All quantities reported are in lattice units.